

A Note on "Exp-function method for the exact solutions of fifth order KdV equation and modified Burgers equation"

Nikolay A. Kudryashov*, Dmitry I. Sinelshchikov

Department of Applied Mathematics, National Research Nuclear University MEPhI, 31 Kashirskoe Shosse, 115409 Moscow, Russian Federation

Abstract

We discuss the recent paper by Inan and Ugurlu [Inan I.E., Ugurlu Y., Exp-function method for the exact solutions of fifth order KdV equation and modified Burgers equation, Appl. Math. Comp. 217 (2010) 1294 – 1299]. We demonstrate that all exact solutions of fifth order KdV equation and modified Burgers equation by Inan and Ugurlu are trivial solutions that are reduced to constants. Moreover, we show exact solutions of the fifth – order equation studied by Inan and Ugurlu cannot be found by the Exp-function method.

Keywords: Nonlinear evolution equations, exact solutions, fifth order KdV equation, modified Burgers equation.

PACS 02.30.Jr - Ordinary differential equations

In the recent paper [1] Inan and Ugurlu considered the fifth – order KdV equation

$$u_t + u u_x + u_{xxxx} = 0 \quad (1)$$

In [1] Inan and Ugurlu used the traveling wave solutions $u(x, t) = u(\xi)$, $\xi = kx - wt$ in Eq.(1) and obtained nonlinear ordinary differential equation in the form

$$w u' + k u u' + k^5 u^{(5)} = 0, \quad (2)$$

where

$$u' = \frac{du}{d\xi}, \quad u^{(5)} = \frac{d^5 u}{d\xi^5}.$$

After integration Eq. (2) with respect to ξ we can have

$$C + w u + \frac{k}{2} u^2 + k^5 u^{(4)} = 0. \quad (3)$$

*nakudryashov@mephi.ru

Inan and Ugurlu omitted the constant of integration C in [1] and essentially reduced a class for exact solutions of Eq.(1). It is known that there are some exact solutions of Eq.(1) expressed through rational and elliptic functions [2–4]. However taking the Exp-function method into consideration we cannot find these solutions.

Using Exp-function method Inan and Ugurlu found the following five solutions of Eq.(1) in [1]

$$u_1(x, t) = \frac{-\frac{2w}{k} \exp(kx + wt) - \frac{2b_0 w}{k} - \frac{2b_{-1} w}{k} \exp(-kx - wt)}{\exp(kx + wt) + b_0 + b_{-1} \exp(-kx - wt)}, \quad (4)$$

$$u_2(x, t) = \frac{-\frac{2w}{k} \exp(kx + wt) - \frac{2b_0 w}{k}}{\exp(kx + wt) + b_0}, \quad (5)$$

$$u_3(x, t) = \frac{-2k^4 \exp(kx + k^5 t) - 2b_0 k^4}{\exp(kx + k^5 t) + b_0}, \quad (6)$$

$$u_4(x, t) = \frac{-32k^4 \exp(kx + 16k^5 t)}{\exp(kx + 16k^5 t)}, \quad (7)$$

$$u_5(x, t) = \frac{-32k^4 \exp(kx + 16k^5 t) - 32b_0 k^4}{\exp(kx + 16k^5 t) + b_0}. \quad (8)$$

However it is easy to note that all these "solutions" can be written as constants

$$u_1(x, t) = \frac{2w}{k}, \quad (9)$$

$$u_2(x, t) = -\frac{2w}{k}, \quad (10)$$

$$u_3(x, t) = -2k^4, \quad (11)$$

$$u_4(x, t) = -32k^4, \quad (12)$$

$$u_5(x, t) = -32k^4. \quad (13)$$

It is wonderful but even trivial "solutions" (9), (11), (12) and (13) do not satisfy Eq.(3) at $C = 0$.

Inan and Ugurlu in [1] also presented the exact solutions of the modified Burgers equation

$$u_t + u^2 u_x + u_{xx} = 0 \quad (14)$$

Authors [1] have used again the traveling wave solutions for Eq. (14). They also have omitted the constant of integration in reduction of (14). Finally, Inan and Ugurlu have considered following equation

$$w u + \frac{k}{3} u^3 + k^2 u' = 0. \quad (15)$$

Equation (15) is well known and was introduced by Jakob Bernoulli in 1695. Eq. (15) can be reduced to the linear form using transformation

$$u(\xi) = \frac{1}{\sqrt{v(\xi)}} \quad (16)$$

In this case Eq.(15) is reduced to the form

$$k^2 v_\xi = \frac{2k}{3} + 2wv \quad (17)$$

The general solution of Eq.(15) can be presented in the form

$$u(\xi) = \pm \frac{3}{\sqrt{9C_2 e^{\frac{2w\xi}{k^2}} - \frac{3k}{w}}}, \quad (18)$$

where C_2 is a arbitrary constant.

In [1] Inan and Ugurlu applied again the Exp-function method and found "three solutions" of Eq. (14). The first "solution" is

$$u_1(x, t) = \frac{\sqrt{\frac{3k}{2}} \exp(kx - \frac{k^2}{2}t) - \sqrt{\frac{3k}{2}}b_0}{\exp(kx - \frac{k^2}{2}t) + b_0} \quad (19)$$

However, solution (19) do not satisfy Eq. (14). Substituting function (19) into Eq. (14) we obtain the following expression

$$E = -\frac{2\sqrt{6}b_0^2 k^{5/2} \exp(kx - \frac{k^2}{2}t) \left(2 \exp(kx - \frac{k^2}{2}t) - b_0\right)}{(\exp(kx - \frac{k^2}{2}t) + b_0)^4} \quad (20)$$

We can see that this expression is not equal to zero in the general case.

The second and the third solutions can be written as

$$u_2(x, t) = \frac{\sqrt{3k} \exp(kx - \frac{k^2}{2}t)}{\exp(kx - \frac{k^2}{2}t)} \quad (21)$$

$$u_3(x, t) = \frac{\sqrt{\frac{3k}{2}} \exp(kx - \frac{k^2}{2}t) + \sqrt{\frac{3k}{2}}b_0 + \sqrt{\frac{3k}{2}}b_{-1} \exp(-kx + \frac{k^2}{2}t)}{\exp(kx - \frac{k^2}{2}t) + b_0 + b_{-1} \exp(-kx + \frac{k^2}{2}t)} \quad (22)$$

However these expressions can be presented as the trivial solutions of Eq.(15). We have

$$u_2(x, t) = \sqrt{3k}, \quad (23)$$

$$u_3(x, t) = \sqrt{\frac{3k}{2}}. \quad (24)$$

However the trivial solutions (23) and (24) do not satisfy Eq.(15) as well.

We can see that all exact solutions obtained by Inan and Ugurlu in [1] are trivial and we can obtain these solutions without the Exp-function method. What is more these trivial solution do not satisfy equations as well.

In conclusion we strongly recommend to authors and referees to look at papers [5–23] carefully before finding exact solutions of nonlinear differential equations.

References

- [1] Inan I.E., Ugurlu Y. Exp-function method for the exact solutions of fifth order KdV equation and modified Burgers equation, Appl. Math. Comp. 217 (2010) 1294 – 1299
- [2] Kudryashov N.A. Analitical theory of nonlinear differential equations, Moskow - Igevsk, Institute of computer investigations, 2004, (in Russian)
- [3] Kudryashov N.A., Nonlinear Differential Equations with Exact Solutions Expressed via the Weierstrass Function, Z. Naturforsch. 59a (2004) 443 454
- [4] Demina M.V., Kudryashov N.A. From Laurent series to exact meromorphic solutions: The Kawahara equation, Physics Letters A 174 (2010) 4023 – 4029
- [5] Kudryashov N.A., Loguinova N.B., Be careful with Exp-function method, Commun. Nonlinear Sci. Numer. Simulat. 2009;14:1881 - 1890
- [6] Kudryashov N.A., On "new travelling wave solutions" of the KdV and the KdV-Burgers equations. Commun Nonlinear Sci Numer Simul 2009;14:1891-900.
- [7] Kudryashov N.A., Seven common errors in finding exact solutions of nonlinear differential equations, Commun Nonlinear Sci Numer Simul 2009;14:35033529.
- [8] Kudryashov N.A., Soukharev M.B. , Popular Ansatz methods and Solitary wave solutions of the Kuramoto-Sivashinsky equation, Regular and Chaotic Dynamics, 14 (2009) 407 - 419
- [9] Kudryashov N.A., Comment on: "A novel approach for solving the Fisher equation using Exp-function method", Physics Letters A 2009;373:1196 - 1197
- [10] Kudryashov N.A. Meromorphic solutions of nonlinear differential equations, Commun Nonlinear Sci Numer Simulat (2009), doi:10.1016/j.cnsns.2009.11.013.
- [11] Kudryashov N.A., Sinelshchikov D.I., A note on "Abundant new exact solutions for the (3+1) – dimensional Jimbo – Miva equation", Journal of Mathematical Analysis and Applications, 371 (1) (2010) 393 – 396
- [12] Kudryashov N.A., Soukharev M.B. Comment on: Multi soliton solution, rational solution of the Boussinesq – Burgers equation, Commun Nonlinear Sci Numer Simul, 15 (2010) 1765 – 1767.
- [13] Kudryashov N.A., A note on New Exact solutions for the Kawahara equation using Exp – function method, Journal of Computational and Applied Mathematics, 234 (12) 3511 – 3512

- [14] Kudryashov N.A., A note on the G'/G – expansion method, *Applied Mathematics and Computation*, 217 (2010) 1755 – 1758
- [15] Kudryashov N.A., A new note on exact complex travelling wave solutions for $(2+1)$ – dimensional B – type Kadomtsev – Petviashvili equation, *Applied Mathematics and Computation*, 217 (2010) 2282 – 2284
- [16] Kudryashov N.A., Sinelshchikov D.I., A Note on the Lie Symmetry Analysis and Exact Solutions for the Extended mKdV Equation, *Acta Applicandae Mathematica*, (2010), doi:10.1007/a10440-010-9582-6
- [17] Parkes E.J., A note on travelling - wave solutions to Lax’s seventh - order KdV equation, *Appl. Math. Comput.* 2009; 215:864 - 865
- [18] Parkes E.J. Observations on the tanh-coth expansion method for finding solutions to nonlinear evolution equations. *Appl. Math. Comput.* 217 (2010), 1749 – 1754
- [19] Parkes E.J., A note on solitary travelling-wave solutions to the transformed reduced Ostrovsky equation, *Commun. Nonlinear. Sci. Numer. Simul.*, 15 (2010), 2769 – 2771
- [20] Parkes E.J., Observations on the basic (G'/G) – expansion method for finding solutions to nonlinear evolution equations, *Appl. Math. Comput.* 217 (2010) 1759 – 1763
- [21] Parkes E.J., A note on "New travelling wave solutions to the Ostrovsky equation" (2010), doi:10.1016/j.amc.2010.08.044
- [22] Popovych R.O., Vaneeva O.O., More common errors in finding exact solutions of nonlinear differential equations, *Commun. Nonlinear. Sci. Numer. Simul.*, 15 (2010), 3887 – 3899
- [23] Sinelshchikov D.I., Comment on: New exact traveling wave solutions of the $(3 + 1)$ -dimensional Kadomtsev-Petviashvili (KP) equation, *Commun. Nonlinear. Sci. Numer. Simul.*, 15 (2010), 3235 – 3236